

	<b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b> FOR INVOICE 162850 DATED 01/01/17	
<b>JAMES T. SMITH DAIRY HEIFER CATTLE FEED</b> <b>MEDICATED</b> TYPE C FEED CONTAINING BOVATEC™(LASALOCID)		
FORMULA #: JTS Heifer 12/16		

**STATEMENT OF PURPOSE**

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). For control of coccidiosis caused by Eimeria bovis and Eimeria zuernii in cattle.

**ACTIVE DRUG INGREDIENT**

Lasalocid(as lasalocid sodium)	32.40 grams/ton 16.20 mg/pound
--------------------------------	-----------------------------------

**USE DIRECTIONS**

**Important:** The lasalocid must be contained in at least 1 pound of feed. If the feeding rate falls less than 1 pound, mix with additional nonmedicated feed so that the drug is contained in at least one pound of feed.

**FOR INCREASED RATE OF WEIGHT GAIN**  
 Feed on a hand-fed basis so that each animal will be consuming between 3.7 and 18.5 pounds per head per day which will provide between 60 to 300 mg lasalocid per head per day when on pasture. Feed continuously.

**NOTE:** Daily intakes of lasalocid in excess of 200 mg/head/day or 12.35 pounds of this feed/head/day, have not been shown to be more effective than 200 mg/head/day.

**FOR COCCIDIOSIS CONTROL**  
 Hand-feed daily at the rate of 1 mg of lasalocid per 2.2 pounds body weight per day to cattle weighing up to 800 pounds with a maximum of 360 mg lasalocid per head per day in accordance with the schedule below. In herds of mixed sizes use creep feeders that prevent animals over 800 pounds access to this feed.

ANIMAL WT. (POUNDS)	LBS TO FEED PER DAY	ANIMAL WT. (POUNDS)	LBS TO FEED PER DAY
100	2.81	500	14.03
200	5.61	600	16.84
300	8.42	700	19.64
400	11.22	800	22.45

**Note:** Coccidiosis may occur when young pasture cattle are co-mingled with adult cattle passing coccidial oocysts.

**PRECAUTIONARY STATEMENTS**

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.  
**CAUTION:** The safety of lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

MANUFACTURED BY:  
 BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

BOVATEC™ is a trademark of Alphaarma Inc. label revision 11/21/2017

	<b>CUSTOMER-FORMULA MEDICATED LABEL ATTACHMENT</b> FOR INVOICE 162850 DATED 01/01/17	
<b>JAMES T. SMITH DAIRY HEIFER CATTLE FEED</b> <b>MEDICATED</b> TYPE C FEED CONTAINING BOVATEC™(LASALOCID)		
FORMULA #: JTS Heifer 12/16		

**STATEMENT OF PURPOSE**

For increased rate of weight gain in pasture cattle (slaughter, stocker, feeder cattle and dairy and beef replacement heifers). For control of coccidiosis caused by Eimeria bovis and Eimeria zuernii in cattle.

**ACTIVE DRUG INGREDIENT**

Lasalocid(as lasalocid sodium)	32.40 grams/ton 16.20 mg/pound
--------------------------------	-----------------------------------

**USE DIRECTIONS**

**Important:** The lasalocid must be contained in at least 1 pound of feed. If the feeding rate falls less than 1 pound, mix with additional nonmedicated feed so that the drug is contained in at least one pound of feed.

**FOR INCREASED RATE OF WEIGHT GAIN**  
 Feed on a hand-fed basis so that each animal will be consuming between 3.7 and 18.5 pounds per head per day which will provide between 60 to 300 mg lasalocid per head per day when on pasture. Feed continuously.

**NOTE:** Daily intakes of lasalocid in excess of 200 mg/head/day or 12.35 pounds of this feed/head/day, have not been shown to be more effective than 200 mg/head/day.

**FOR COCCIDIOSIS CONTROL**  
 Hand-feed daily at the rate of 1 mg of lasalocid per 2.2 pounds body weight per day to cattle weighing up to 800 pounds with a maximum of 360 mg lasalocid per head per day in accordance with the schedule below. In herds of mixed sizes use creep feeders that prevent animals over 800 pounds access to this feed.

ANIMAL WT. (POUNDS)	LBS TO FEED PER DAY	ANIMAL WT. (POUNDS)	LBS TO FEED PER DAY
100	2.81	500	14.03
200	5.61	600	16.84
300	8.42	700	19.64
400	11.22	800	22.45

**Note:** Coccidiosis may occur when young pasture cattle are co-mingled with adult cattle passing coccidial oocysts.

**PRECAUTIONARY STATEMENTS**

**WARNING:** A withdrawal period has not been established for this product in pre-ruminating cattle. Do not use in calves to be processed for veal.  
**CAUTION:** The safety of lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

MANUFACTURED BY:  
 BLUE BIRD FEED MILL, ANY CITY, ANY STATE 55555

Net Weight on Bag and/or Invoice

BOVATEC™ is a trademark of Alphaarma Inc. label revision 11/21/2017